

Thermal decoupling of WIMPs from first principles

Torsten Bringmann

SISSA/ISAS, via Beirut 2-4, 34013 Trieste, Italy

E-mail: bringman@sissa.it

Stefan Hofmann

Perimeter Institute for Theoretical Physics, Waterloo, Ontario, N2L2Y5, Canada

E-mail: shofmann@perimeterinstitute.ca

Abstract. Weakly interacting massive particles (WIMPs) are arguably the most natural DM candidates from a particle physics point of view. After their number density has frozen out in the early universe, thereby setting their relic density today, WIMPs are still kept very close to thermal equilibrium by scattering events with standard model particles. The complete decoupling from the thermal bath happens as late as at temperatures of around $\sim 1 - 10$ MeV and provides an important cosmological scale that can directly be translated into a small-scale cutoff of the spectrum of matter density fluctuations. We present here a full analytic treatment of the kinetic decoupling process from first principles. This allows an exact determination of the decoupling scale, for arbitrary WIMP candidates and any scattering amplitude. As an application, we consider the situation of the lightest supersymmetric particle as well as the lightest Kaluza-Klein particle that arises in theories with universal extra dimensions; furthermore, we show that our formalism can also easily be applied to, e.g., the evolution of the non-relativistic electrons into the recombination regime. Finally, we comment on the impacts for the smallest gravitationally bound structures and the prospects for the indirect detection of dark matter.

1. Introduction

Based on the simplest inflationary scenario, a flat cosmological model with dark energy and cold dark matter as dominant contingents in the cosmic energy budget fits the spectrum of observed temperature anisotropies imprinted in the cosmic microwave background radiation (CMB) [1]. Assuming that the primordial fluctuations are adiabatic with a power law spectrum, the temperature anisotropy data *alone* indicates a significant amount of non-baryonic cold dark matter (CDM); analysis of the anisotropies finds that the relative matter density $\omega_m \equiv \Omega_m h^2 = 0.127^{+0.007}_{-0.013}$ is roughly six times larger than the relative baryon density. This is consistent with a diversity of other cosmological observations, e.g. the analysis of galaxy clustering as observed by SDSS [2] and 2dFGRS [3], improvements in the lensing data [4], as well as improvements in small-scale CMB measurements, e.g. [5].

Arguably the best studied class of CDM particle candidates are weakly interacting massive particles, or WIMPs (for reviews on particle DM, see [6]). Equipped with masses and typical interaction strengths set by the electroweak scale, they naturally occur in many extensions to the standard model of particle physics (SM), e.g. in its supersymmetric versions or in models with space-like extra dimensions. Thermally produced in the early universe, and provided their stability, WIMPs automatically acquire a relic abundance that is of the right order of magnitude to account for the observed matter density. Another well-studied CDM particle candidate is the axion. The essential difference between WIMPs and axions is that the former have been kept in local thermal equilibrium prior to some epoch in the cosmological evolution while the latter have not. This is because by the time the axion potential developed a minimum at around the QCD phase transition, and the axions thus became massive, any interactions were already heavily suppressed by the Peccei-Quinn scale. In this article, we will focus solely on WIMPs and often use the abbreviation CDM as a synonym for these.

It has long been a widely held belief that the process of structure formation is completely independent of the CDM particle nature; CDM density perturbations, in particular, should only be effected by their gravitational interaction and bear no visible imprint of particle interactions. In this case, however, cosmological perturbation theory predicts a monotonously growing support for structure formation on ever smaller scales. More precisely, the spectrum of matter density perturbations grows logarithmically with the comoving wavenumber that characterizes the perturbation mode [7, 8]. This behaviour has recently been confirmed yet another time, down to the resolution of a numerical simulation with unprecedented accuracy [14]. As a consequence, a regularization scheme would be required to render linear perturbation theory well-defined, e.g. a cut-off in the wavenumber that corresponds to some resolution limit. From cosmological perturbation theory we know that gravitational interactions alone cannot give rise to a fundamental cut-off scale, so it has to be introduced by hand. On the other hand, it is of course rather problematic to impose such an ad hoc small-scale cut-off since, in hierarchical structure formation, small scale perturbations enter the

nonlinear regime first and structures on larger scales form only later via mergers and accretion.

In Ref. [9] it was shown that CDM interactions with the radiation component give rise to a natural cut-off that can be calculated directly from the interaction theory underlying a specific dark matter candidate. The authors of [10] confirmed the results of [9]. The (physical) cut-off is set by viscous processes around the kinetic decoupling of CDM from radiation. Close to kinetic decoupling, CDM density perturbations receive in-medium modifications that give rise to a dispersion relation with a non-vanishing imaginary contribution and thus to absorption. Subsequently, free streaming of CDM suppresses further the power of structure formation on small scales. In that way, collisional and collision-less damping processes give rise to a fundamental cut-off in the spectrum of linear CDM perturbations [7, 8] and thus to a characteristic feature that clearly depends on the CDM micro-physics. It was later shown numerically [11] that acoustic oscillations have to be taken into account at kinetic decoupling, however, the order of magnitude of the cut-off scale is correctly given in [7, 8]. Most recently, the author of [12] succeeded in including acoustic oscillations before and during kinetic decoupling, the viscous coupling during kinetic decoupling and free streaming in a consistent kinetic treatment. Close to this physical cut-off scale, the power spectrum develops a maximum. Hence, this scale characterizes those density perturbations that typically enter the non-linear regime first and therefore give rise to the first purely gravitationally bound structures.

Provided that the dark matter density distribution today retains traces of these typical first halos, the cut-off scale described above may thus have important observational consequences. While it is certainly true that the present CDM density profile is independent of the particle nature of CDM on super-kpc scales, this is highly questionable on sub-pc scales. Let us therefore stress that WIMP detection experiments actually probe the dark matter density profile on sub-mpc scales [13]: for a given WIMP candidate, the prospects for a direct detection (via elastic scattering processes with the detector nuclei) depend on the effective reaction volume which is given by the detector size and the path the detector travels during data taking. Indirect detection experiments, on the other hand, aim at measuring flux densities from dark matter annihilation products, which depend quadratically on the dark matter density contrast in the observed region. Substructures thus enhance the fluxes and therefore the prospects for indirect detection, probing the dark matter density profile on sub-galactic scales.

Let us stress that it is, in fact, the temperature at kinetic decoupling that provides the interface, at least in principal, between the interaction theory underlying a specific CDM candidate and CDM substructures that are probed by direct and indirect dark matter searches. Motivated by this observation, we present in this article a complete and analytic description of the kinetic decoupling of WIMPs from the local heat bath, without assuming a specific WIMP candidate. In Section 2, we introduce important time scales that characterize the non-equilibrium state of WIMPs and discuss on a very

qualitative level both chemical and kinetic decoupling of WIMPs, before we turn to the presentation of a precise treatment of the kinetic decoupling process. We introduce a generalized WIMP temperature (which characterizes the WIMP distribution even in the kinetic stage and agrees with the heat bath temperature before kinetic decoupling) and derive a master equation for the evolution of this temperature. Section 3 is devoted to various examples: as a first and familiar example we quantify, for illustration, the deviation of the electron temperature from the photon temperature during the recombination regime. More relevant for our original purposes is the next example, the kinetic decoupling of bino-like CDM from SM fermions. As a last example we consider Kaluza-Klein dark matter. Section 4 provides a summary of the processes that constitute the small-scale cut-off in the linear matter power spectrum; here, we also shortly review what is known about the total mass fraction in and survival probability of the smallest gravitationally bound structures that correspond to this scale. In section 5 we discuss prospects for indirect dark matter detection experiments in light of small-scale structure formation. Section 6, finally, concludes. All technical details and derivations are presented in the three appendices: In Appendix A, we solve the Boltzmann equation for the collision integral, as determined in Appendix B, that describes the scattering processes of relevance to our discussion; in Appendix C, we present the LKP scattering amplitude and use this opportunity to demonstrate the applicability of our formalism even in the case of s -channel resonances of the scattering amplitude.

2. Thermal relics and their decoupling from the heat bath

In this section, we introduce various time scales that allow to characterize different stages in the evolution of WIMPs. While a thorough understanding of the various characteristic time scales is not required for the later computation of the kinetic decoupling temperature, the presentation mainly aims at clarifying the importance of the relaxation time for WIMPs close to local thermal equilibrium. We then discuss chemical and kinetic decoupling of WIMPs from the local heat bath on a very qualitative level, before proceeding to a precise description of kinetic decoupling – which is the main concern of this article.

2.1. Characteristic time scales

The most detailed statistical description of WIMPs is given by the N -particle phase space density, with N denoting their number in a comoving Volume V . Around kinetic decoupling, the interaction radius r_{int} of WIMPs (defined as the maximal distance between a WIMP and SM – or, rather, heat bath – particles over which interactions can occur) is much smaller than the mean distance between the particles, i.e. $Nr_{\text{int}}^3/V \ll 1$. This simple inequality allows the introduction of different time scales that are important for the classical kinetic description of dark matter. The first time scale is called *collision time* and defined by $\tau_c \equiv r_{\text{int}}/\langle v \rangle$, with $\langle v \rangle$ denoting the mean dark matter

velocity. A second time scale is provided by the time between two subsequent collisions, $\tau_f \equiv l_f/\langle v \rangle$, where l_f is the mean free path of dark matter particles. The third and for our considerations most important time scale is the *relaxation time* τ_r . It marks the time during which local thermal equilibrium is established in a comoving volume that contains many dark matter particles, but that is still small compared to V . The last time scale we want to consider here, τ_{eq} , is given by the time during which global equilibrium in V is established.

From elementary kinetic theory it follows that the mean free path can be approximated by $l_f/V^{1/3} \sim (1/N)(V^{1/3}/r_{int})^2$, hence $\tau_c \ll \tau_f$. Since CDM particles with weak interactions need many collisions in order to maintain local thermal equilibrium, we also have $\tau_f \ll \tau_r$. Finally, one evidently has $\tau_r \ll \tau_{eq}$. To summarize, we get the following chain of inequalities for the characteristic time scales introduced above:

$$\tau_c \ll \tau_f \ll \tau_r \ll \tau_{eq}. \quad (1)$$

For a description of processes on time scales $\delta t \ll \tau_c$, the full N -particle phase space density is required, and a reduced description in terms of some $\tilde{N} < N$ -particle phase space distribution is impossible. A reduced kinetic description is, however, available for processes with $\tau_c \ll \delta t \ll \tau_r$. The WIMPs can then adequately be described by a one-particle distribution that determines the probability distribution for the locations and the momenta of the CDM particles. For processes on time scales $\tau_r \ll \delta t \ll \tau_{eq}$, the system is in local thermal equilibrium and further simplifications apply. During this stage, the WIMPs are fully described by the local number, momentum and kinetic energy density, which are given as the zeroth, first and second moments, respectively, of the one-particle distribution with respect to the WIMP momentum. This is the underlying reason that allows to describe such a system as a perfect or almost perfect fluid.

2.2. Scales of decoupling

Initially, the WIMP system is (almost) in local chemical and thermal equilibrium, with chemical equilibrium provided by the detailed balance between WIMP annihilation and production processes, and local thermal equilibrium maintained by elastic scattering processes with SM particles in the heat bath. Typically, chemical and thermal equilibrium are protected by the aforementioned processes for temperatures above the WIMP particle mass, $T \gtrsim M_\chi$.

As the Universe expands, the temperature decreases and the WIMP momenta become redshifted. At $T \sim M_\chi$ detailed balance is destroyed: the lighter species, dominating the energy budget, can no longer produce WIMPs, while the reverse channel is still open. As a consequence, between the end of detailed balance at $T \sim M_\chi$ and chemical decoupling at $T = T_{cd}$, the total (comoving) number of WIMPs decreases. Chemical decoupling happens roughly when the WIMP annihilation rate drops below the Hubble rate H , and, as a consequence, annihilation processes cease to proceed and the relic dark matter abundance is fixed. This is expected to happen since the relevant target

density in the annihilation processes is provided by the WIMP number density, which is Boltzmann suppressed for temperatures $T < M_\chi$. For instance, in the case of WIMPs the chemical decoupling scale is typically given by $T_{\text{cd}} \sim M_\chi/25$. Hence, the total number of WIMPs is conserved for $T \gtrsim M_\chi$ – dynamically protected by detailed balance, decreases during $M_\chi/25 \lesssim T \lesssim M_\chi$, and again protected for $T \lesssim M_\chi/25$ – basically by the volume expansion.‡ Note that the dependence of the chemical decoupling scale T_{cd} on the WIMP annihilation rate (and thus on the underlying interaction theory) can be completely eliminated by using the value for the relative WIMP density as inferred e.g. from the temperature anisotropy data.

After chemical decoupling elastic scattering processes still keep WIMPs in local thermal equilibrium. It is very important to realize that even though dark matter is now a chemically distinct particle species, it is still a constituent of the local heat bath. The relevant target density for elastic scattering processes is provided by the number density of relativistic SM particles and thus only decreasing as T^3 . Eventually, also the elastic scattering rate Γ_{el} drops below the Hubble rate and elastic scattering processes cease. However, even before last scattering, local thermal equilibrium cannot be maintained since the relaxation processes within the WIMP system are weakened by the Hubble expansion. Consequently, the relevant time scales are the relaxation time τ_r (intrinsic) and the Hubble time H^{-1} (extrinsic).

Before we will present a way to analytically determine the decoupling scale in the next Section, let us now roughly estimate the relaxation time for WIMPs as follows (see also [9]): our system consists of a weakly interacting particle species with mass M_χ and a local heat bath characterized by a temperature $T \ll M_\chi$, so M_χ is the dominant energy scale. The average (three-) momentum transfer Δp from a light (heat bath) particle to the WIMP, during a single elastic scattering process, is thus roughly given by its energy, i.e. $\Delta p \sim T$. We assume now that Δp is constant during the relaxation process. After N_{coll} collisions, inducing a random walk in momentum space, the WIMP momentum will thus have changed by $\Delta p (N_{\text{coll}})^{1/2}$. In order to maintain local thermal equilibrium, this should be comparable to the typical momentum for thermally distributed WIMPs, which is given by $(M_\chi T)^{1/2}$. The number of collisions required is thus $N_{\text{coll}} \sim M_\chi/T \gg 1$ and the relaxation time becomes $\tau_r \sim N_{\text{coll}}/\Gamma_{\text{el}} \sim (m/T)1/\sigma_{\text{scale}}$. The scale for the total elastic scattering process can furthermore be assumed to be approximated by $\sigma_{\text{scale}} \equiv (G_F m_W^2) M_\chi^2/m_Z^4$, with G_F denoting Fermi's constant, m_W and m_Z are the masses of the charged and neutral weak gauge bosons, respectively. Kinetic decoupling occurs approximately at the temperature T_{kd} , for which $\tau_r(T_{\text{kd}}) = H^{-1}(T_{\text{kd}})$. With these simplifications, we thus expect $T_{\text{kd}} \sim M_\chi^{1/4} m_Z/M_{\text{Pl}}^{1/4} (G_F m_W^2)^{1/2} \sim \mathcal{O}(1-10)$ MeV. Let us stress again that this is only a rough estimate for the decoupling scale; below, we will determine the precise answer by solving the appropriate kinetic theory.

‡ Strictly speaking, the relic abundance is protected only until density perturbations generate density contrasts that again allow for WIMP annihilations.

2.3. A precise description of the kinetic decoupling process

Around kinetic decoupling WIMPs pass from the hydrodynamical to the kinetic stage. Their appropriate description changes accordingly: while the hydrodynamical stage describes the evolution of fluid variables that are given as expectation values with respect to the one-particle phase space distribution, the kinetic stage is concerned with the evolution of the one-particle phase space distribution $f(\mathbf{p})$ itself.

A more precise determination of the decoupling temperature requires the introduction of a temperature parameter T_χ for dark matter, which we choose to define as the second moment of WIMP phase-space distribution function f :

$$T_\chi \equiv \frac{2}{3} \left\langle \frac{\mathbf{p}^2}{2M_\chi} \right\rangle, \quad \text{with } \langle \mathcal{O}(\mathbf{p}) \rangle \equiv \frac{1}{n_\chi} \int \frac{d^3p}{(2\pi)^3} \mathcal{O}(\mathbf{p}) f(\mathbf{p}). \quad (2)$$

Here, n_χ denotes the WIMP number density and $\mathcal{O}$ is some microscopic observable. The parameter T_χ that we introduced above agrees with the heat bath temperature for $T > T_{\text{kd}}$ and can be viewed as a generalized WIMP temperature for $T < T_{\text{kd}}$; its difference to T thus characterizes the deviation of WIMPs from the equilibrium state.

The dynamics of the one-particle phase space distribution, governed by the Boltzmann equation, is nontrivial during the kinetic decoupling process (kinetic stage). In the case of dark matter weakly coupled to radiation the main complication arises from the coupling of slow processes (non-relativistic propagation of dark matter) and fast processes (relativistic propagation of radiation) in the collision integral, which is enforced via the explicit energy momentum conservation constraining the phase space available for elastic scattering processes.

Around kinetic decoupling, the WIMP mass M_χ is by far the largest energy scale involved. In particular, the kinetic WIMP energy, as well as the average momentum transfer during the elastic scattering processes, is tiny compared to M_χ . This allows to factorize slow and fast processes in the collision integral by utilizing a Born-Oppenheimer type of approximation and expanding the collision integral in $\mathbf{p}^2/M_\chi^2$. Instead of solving the Boltzmann equation for f , we can then solve its second moment and derive, in Appendix A, the following process equation for $T_\chi(T)$:

$$\frac{dT_\chi}{dT} - \left[2 + a \left(\frac{T}{M_\chi} \right)^{n+2} \right] \frac{T_\chi}{T} = -a \left(\frac{T}{M_\chi} \right)^{n+2}. \quad (3)$$

Here, n is determined by the low-energy behaviour of the scattering amplitude at zero momentum transfer, $|\mathcal{M}|_{t=0}^2 \propto (\omega/M_\chi)^n$, where ω is the energy of the SM scattering partner; a is a numerical constant that is defined in (A.23). The uniquely defined solution to Eq. (3) with the correct asymptotic behavior in the high-temperature limit, $T_\chi \xrightarrow{T \rightarrow \infty} T$, is given by:

$$T_\chi = T \left\{ 1 - \frac{z^{1/(n+2)}}{n+2} \exp[z] \Gamma[-(n+2)^{-1}, z] \right\}_{z=\frac{a}{n+2} \left(\frac{T}{M_\chi} \right)^{n+2}}. \quad (4)$$

In the low-temperature limit, $T \rightarrow 0$, we have $T_\chi \sim T^2/M_\chi$. Since, as we shall see, kinetic decoupling occurs on a rather short time scale, a suitable choice for a *definition*

of the kinetic decoupling temperature is obtained by matching the two above-mentioned asymptotic solutions to Eq. (3), resulting in:

$$\frac{T_{\text{kd}}}{M_\chi} \equiv \left(\frac{T^2}{M_\chi T_\chi} \right)_{T \rightarrow 0} = \left(\left(\frac{a}{n+2} \right)^{1/(n+2)} \Gamma \left[\frac{n+1}{n+2} \right] \right)^{-1}. \quad (5)$$

For a more detailed discussion of the process equation (3), we refer to Appendix A and the following Section, where we consider some examples of particular interest and determine the exact values of the decoupling temperature in these cases. However, from (A.23), let us already now remark that we generically expect it to be of the order of

$$\frac{T_{\text{kd}}}{M_\chi} \sim (\alpha^{-2} M_\chi / M_{\text{Pl}})^{1/(n+2)}, \quad (6)$$

where α is the effective coupling constant that appears in the scattering amplitude. For example, we have $\alpha = \alpha_{\text{em}}$, $n = 0$ in the case of recombination (Thompson scattering) and $\alpha = \alpha_Y$, $n = 2$ for the thermal decoupling of Binos or Kaluza-Klein photons. From Eq. (6) we can see that, indeed, the WIMP micro-physics enters in a crucial way in the determination of the kinetic decoupling scale and thus – see Section 4 – in the numerical value for the cutoff in the spectrum of CDM density perturbations.

3. Examples

In this section, we apply the formalism developed in Appendix A and B, as reviewed in the preceding Section, for illustrative purposes first to the decoupling of electrons from radiation during the recombination era. In a second step, we turn closer to the main focus of this article and discuss the kinetic decoupling of typical WIMP DM candidates, i.e. bino-like and Kaluza-Klein dark matter, in particular.

3.1. Recombination

After the (chemical) decoupling of electrons and positrons at a temperature of around $T \sim 1$ MeV, and the subsequent annihilation of e^+e^- pairs, the photons were still interacting with protons and the remaining electrons, mainly through Compton scattering with the latter. Such a plasma of interacting relativistic and non-relativistic particles cannot, strictly speaking, be kept in thermal equilibrium in an expanding universe. On the other hand, CMB observations show a black-body spectrum to an accuracy of about 10^{-4} [15]. As a first application of our formalism, we will now quantify the expected deviation from a thermal distribution at the time when the universe became transparent and the CMB photons that we observe today were released. For a similar treatment of the transition into the recombination regime, see, e.g. [16].

At zero momentum transfer and for small fermion energies, the amplitude for Thompson scattering is given by

$$|\mathcal{M}|_{t=0}^2 = 64\pi^2 \alpha_{\text{em}}^2. \quad (7)$$

From our general solution (4), the temperature of the non-relativistic electrons therefore evolves as

$$T_e = T \left\{ 1 - \frac{z^{1/2}}{2} \exp[z] \Gamma[-1/2, z] \right\}_{z=\frac{a}{2} \left(\frac{T}{m_e}\right)^2} \quad (8)$$

$$= T \sqrt{\pi z} \exp[z] \{1 - \text{Erf}[\sqrt{z}]\}_{z=\frac{a}{2} \left(\frac{T}{m_e}\right)^2}, \quad (9)$$

where

$$a \stackrel{\text{(A.23)}}{=} \sqrt{\frac{10}{(2\pi)^9 \cdot 3.36}} 128 \pi^2 \alpha_{\text{em}}^2 N_3^- \frac{M_{\text{Pl}}}{m_e} = 1.8 \times 10^{19}. \quad (10)$$

We thus find that when entering the recombination regime at around $T \sim 0.3 \text{ eV}$, the electron temperature deviates only by $(T - T_e)/T = 1.6 \times 10^{-7}$ from the photon temperature. The electrons, in other words, do not in any significant way affect the photon temperature and the photons thus follow an equilibrium distribution into the recombination regime, explaining the observed black-body nature of the CMB. This argument can be made even more quantitative by considering the Boltzmann equation for the photons which, in this regime, is known as the Kompaneets equation [17]:

$$(\partial_t - H\omega\partial_\omega) g(\omega, t) = \frac{8\pi\alpha_{\text{em}}^2}{3m_e^3} n_e \frac{1}{\omega^2} \partial_\omega \left[\omega^4 \left\{ T_e \partial_\omega g(\omega, t) + [1 + g(\omega, t)] g(\omega, t) \right\} \right], \quad (11)$$

where g and ω are the photon distribution function and energy, respectively. One may now *define* the photon temperature by the relation

$$g(1 + g) = -T\partial_\omega g, \quad (12)$$

which is automatically satisfied for a thermal distribution. With this definition, one realizes that for $T = T_e$ the collision term on the right hand side of (11) vanishes identically and the photons therefore just follow an equilibrium distribution.

3.2. Neutralino dark matter

One of the most well-motivated and extensively studied DM candidates is the neutralino. In most supersymmetric extensions to the SM, it appears as the lightest supersymmetric particle (LSP) and is given by a linear combination of the super-partners of the gauge and Higgs fields,

$$\chi \equiv \tilde{\chi}_1^0 = N_{11} \tilde{B} + N_{12} \tilde{W}^3 + N_{13} \tilde{H}_1^0 + N_{14} \tilde{H}_2^0. \quad (13)$$

Here, in order to illustrate our formalism and to describe the kinetic decoupling process for a typical WIMP dark matter candidate, we consider the simple case of a Bino-like neutralino, $\chi \sim \tilde{B}$. From the WMAP relic density constraint, Binors have a mass of a few hundred GeV or less.

To lowest order in the fermion energy ω , the scattering amplitude of Binors with SM fermions is given by [9]:

$$|\mathcal{M}|_{t=0}^2 = 32g_Y^4 (Y_d^4 + Y_s^4) \left(\frac{M_{\tilde{B}}}{M_L^2 - M_{\tilde{B}}^2} \right)^2 \omega^2. \quad (14)$$




Figure 1. The LSP temperature T_χ as a function of the background temperature T in the early universe is shown as a solid line, for the case of a Bino with $M_{\tilde{B}} = 100$ GeV and $M_{\tilde{L}} = 150$ GeV. The rather abrupt transition to a regime with $T_\chi \propto T^2$, where the LSP is completely decoupled from the thermal bath, makes it straightforward to identify the temperature T_{kd} of thermal decoupling as given in Eq. (18); here, it is indicated by a dashed line. For comparison, the (green and blue) dotted lines show the naive estimates for the evolution of T_χ , assuming a sudden decoupling from the thermal bath at the time of last scattering or the relaxation time, respectively (see also [9]).

From (4), we thus find that the neutralino temperature evolves with the background temperature as

$$T_\chi/T = \left\{ 1 - \frac{z^{1/4}}{4} \exp[z] \Gamma[-1/4, z] \right\}_{z=\frac{a}{4}(T/M_{\tilde{B}})^4}, \quad (15)$$

where

$$a \stackrel{\text{(A.23)}}{=} \frac{31}{21} \sqrt{\frac{5\pi^3}{g_{\text{eff}}}} \frac{M_{\text{Pl}}}{M_{\tilde{B}}} g_Y^4 \sum g_{\text{SM}}(Y_d^4 + Y_s^4) \left(\frac{M_{\tilde{B}}^2}{M_{\tilde{L}}^2 - M_{\tilde{B}}^2} \right)^2 \quad (16)$$

$$= 5.5 \times 10^{16} \left(\frac{M_{\tilde{B}}^2}{M_{\tilde{L}}^2 - M_{\tilde{B}}^2} \right)^2 \left(\frac{M_{\tilde{B}}}{100 \text{ GeV}} \right)^{-1}. \quad (17)$$

This corresponds to a decoupling temperature of

$$T_{\text{kd}} \stackrel{(5)}{=} 7.5 \times \left(\frac{M_{\tilde{L}}^2}{M_{\tilde{B}}^2} - 1 \right)^{1/2} \left(\frac{M_{\tilde{B}}}{100 \text{ GeV}} \right)^{5/4} \text{ MeV}. \quad (18)$$

When computing the sum over all contributions, we have assumed an identical mass for all relevant sleptons $\tilde{L}$ (i.e. (anti-)selectrons and (anti-)sneutrinos).

In Fig. 1, we show the evolution of the CDM temperature for a typical Bino DM candidate during the process of kinetic decoupling. For reference, we also indicate the naive estimates for the decoupling temperature that were reviewed in Section 2. At

temperatures only slightly greater than T_{kd} , the LSP is still very tightly bound to the background temperature T (for reference, see also Fig. A1). We therefore expect our result to be highly independent of the details of the QCD phase transition at around $T_{\text{QCD}} \sim 150$ MeV. A second important observation is the rather sharp turnover from $T_\chi \approx T$ to a regime with $T_\chi \propto T^2$, where the LSP has completely decoupled from the thermal bath and simply cools down due to the Hubble expansion. Finally, to take up our previous discussion on the various scales that are involved in the decoupling process, we can clearly see that the relevant quantity is not the time of last scattering, which happens long after the LSP has already decoupled. The relaxation time τ_{rel} , on the other hand, gives as expected a much better estimate for the time of decoupling. The reason that it *underestimates* the actual decoupling time is to be found in the fact that close to decoupling only small deviations from a thermal distribution need to be restored – while the derivation of the relaxation time τ_r implicitly assumed that thermal equilibrium had to be restored from a highly non-thermal distribution.

Obviously, our formalism can in principle easily be extended beyond the simple Bino case that we have presented here and applied to an arbitrary neutralino candidate. In particular, it would be interesting to perform a scan over, e.g., the MSSM parameter space to better understand the expected spread of the decoupling temperature. We note that a first step into this direction has been performed in Ref. [18], based on the estimate that the relaxation time as defined before is a good approximation for the decoupling time; in the case of the Bino, this would correspond to a kinetic decoupling temperature of [9]

$$T_{\text{kd}}^{\text{old}} \sim 20 \times \left(\frac{M_{\tilde{L}}^2}{M_{\tilde{B}}^2} - 1 \right)^{1/2} \left(\frac{M_{\tilde{B}}}{100 \text{ GeV}} \right)^{5/4} \text{ MeV}, \quad (19)$$

which we have, for comparison, also included in Fig. 1. We would like to stress that it would be warranting to re-evaluate the results of Ref. [18] in light of the method that we have presented here; in particular, since we are now able to actually follow the details of the kinetic decoupling process, the underlying uncertainty in the determination of the cutoff-scale is greatly reduced. Also, the method can be applied to regions of the parameter space where a simple scaling of the cross section with energy, like $\sigma \sim \text{const.}$ or $\sigma \propto E_t^2$, is not valid anymore (see Appendix C).

3.3. Kaluza-Klein dark matter

Models with universal extra dimensions (UED) [19], where all standard model (SM) fields are allowed to propagate in a higher-dimensional bulk, have received a great deal of attention since it was realized that they naturally give rise to a new class of dark matter candidates [20, 21]: the higher-dimensional extra degrees of freedom appear in the low-energy effective 4D theory as towers of new, heavy states, the lightest of which – similar to the case of R -parity in supersymmetry – is stable due to an internal Z_2 symmetry (“KK-parity”) that appears as a remnant of the higher-dimensional

translational invariance; thermally produced in the early universe, the lightest Kaluza-Klein particle (LKP) acquires the right WMAP relic density for a compactification scale of about $R^{-1} \sim 1 \text{ TeV}$ [22, 23, 24].

At tree-level, all Kaluza-Klein modes are essentially degenerate in mass, $M \approx R^{-1}$, so radiative corrections become important for the determination of the actual mass hierarchy between these states and, in particular, for the determination of the LKP. Unfortunately, the radiative mass shifts depend in principle on the (unknown) physics at the cutoff scale Λ and should in that sense be treated as free – though small – parameters. However, one usually follows the (self-consistent) assumption that the corresponding contributions at that scale are negligible [20]. In this minimal UED (mUED) model, all masses are determined by only two parameters, R and Λ , and the LKP turns out to be the $B^{(1)}$, the first KK excitation of the weak hypercharge gauge boson.

In Appendix C, we calculate the scattering amplitude for the LKP with SM fermions; to lowest order in the fermion energy ω , it is given by :

$$|\mathcal{M}|_{t=0}^2 = \frac{16}{3} g_Y^4 \left\{ Y_d^4 \left(\frac{M_{B^{(1)}}}{M_{f_d^{(1)}}^2 - M_{B^{(1)}}^2} \right)^2 + Y_s^4 \left(\frac{M_{B^{(1)}}}{M_{f_s^{(1)}}^2 - M_{B^{(1)}}^2} \right)^2 \right\} \omega^2 \quad (20)$$

$$\approx \frac{4}{3} g_Y^4 Y_s^4 \delta_s^{-2} \omega^2, \quad (21)$$

where in the last step we used the fact that the fermion singlet KK states receive much smaller radiative corrections to their masses than the KK doublet states and introduced

$$\delta_s \equiv (M_{f_s^{(1)}} - M_{B^{(1)}})/M_{B^{(1)}} \ll 1. \quad (22)$$

We note that the amplitude takes the same form as for a Bino LSP. The evolution of the LKP temperature is thus described by Eq. (15), but now with

$$a = \frac{31}{126} \sqrt{\frac{5\pi^3}{g_{\text{eff}}}} \frac{M_{\text{Pl}}}{M_{B^{(1)}}} g_Y^4 \sum g_{\text{SM}} \left\{ Y_d^4 \left(\frac{M_{B^{(1)}}}{M_{f_d^{(1)}}^2 - M_{B^{(1)}}^2} \right)^2 + Y_s^4 \left(\frac{M_{B^{(1)}}}{M_{f_s^{(1)}}^2 - M_{B^{(1)}}^2} \right)^2 \right\} \quad (23)$$

$$\approx 2 \times 10^{14} \delta_s^{-2} \left(\frac{M_{B^{(1)}}}{\text{TeV}} \right)^{-1}. \quad (24)$$

The corresponding decoupling temperature is therefore given by

$$T_{\text{kd}} \stackrel{(5)}{\approx} 3 \times 10^2 \delta_s^{1/2} \left(\frac{M_{B^{(1)}}}{\text{TeV}} \right)^{5/4} \text{ MeV}. \quad (25)$$

The sum that appears above runs over all SM fermions that contribute to the scattering process; for temperatures around the kinetic decoupling, these are only electrons and the three neutrino species, as well as the corresponding antiparticles (the contribution from all other particles is strongly Boltzmann suppressed). The mass-splitting between the LKP and the corresponding KK states is then given by $\delta \sim 10^{-2}$. In Figure 2, we take the exact mUED values for δ and show T_{kd} as a function of R and Λ , using the full




Figure 2. The left part of this figure shows the kinetic decoupling temperature T_{kd} for the LKP in the mUED model. From top to bottom, the cutoff scale is set to $\Lambda R = 40, 30, 20$ and the grey band shows the region consistent with the 2σ WMAP relic density constraint for a Higgs mass $m_h \lesssim 150$ GeV (for higher Higgs masses, the grey region broadens and shifts to the right, up to $R^{-1} \lesssim 1.3$ TeV [24]). In the right panel, we leave the mUED setup and treat the mass splitting between the $B^{(1)}$ and the KK leptons as a single, free parameter δ ; from bottom to top, the curves correspond to $\delta = 0.01, 0.05, 0.2$.

expression (23). We find that the LKP decoupling temperature falls into a very narrow range,

$$27 \text{ MeV} \lesssim T_{\text{kd}} \lesssim 34 \text{ MeV}, \quad (26)$$

depending on the parameters of the model. The low value of T_{kd} , despite a TeV scale LKP mass, justifies the above statement about the number of relativistic SM degrees of freedom close to decoupling. In particular, we would like to stress that the kinetic decoupling of the LKP – just as for the Bino – happens sufficiently late so as not to be influenced by the details of the QCD phase transition at around $T_{\text{QCD}} \sim 150$ MeV.

As a next step, we relax the assumptions of the mUED setup and treat the LKP mass and the mass splitting as independent parameters that may be varied about the mUED values. As can be seen from the right panel of Fig. 2, one may in this way encounter higher decoupling temperatures – but without invoking extreme fine-tuning, one will still stay below $T_{\text{kd}} \lesssim 100$ MeV.

4. From kinetic decoupling to damping scales in the power spectrum

After having provided a detailed description of the kinetic decoupling process in the preceding part of this article, we would now like to briefly discuss in the next two Sections, on a rather qualitative level, the further evolution of the WIMP population in the expanding universe.

To start with, and in order to get a qualitative intuition for the physical processes leading to kinetic damping of CDM perturbations, let us use the hydrodynamical description for discussing the evolution of these perturbations. For temperatures $T > T_{\text{kd}}$, i.e. before kinetic decoupling, small-scale perturbations in the CDM fluid are damped due to the tight coupling between the CDM and radiation fluid. Note that this

includes temperatures $T < M_\chi$, when the CDM fluid is already non-relativistic. Around kinetic decoupling, $T \sim T_{\text{kd}}$, the CDM fluid starts to deviate from an ideal fluid, i.e. the coupling to the radiation component is not strong enough to maintain local thermal equilibrium for the CDM any longer. However, the residual coupling between these two fluids generates some friction between them. The main viscous processes causing this friction are bulk- and shear-viscosity, while heat conduction is sub-dominant. Hence, perturbations of the CDM energy momentum tensor behave like sound waves propagating through an opaque medium (the heat bath) [25]. Their dispersion relation receives in-medium modifications, resulting in an imaginary frequency that is generated by the viscous coupling between the CDM perturbations and the heat bath.

The strength of the viscous couplings can not be calculated within the hydrodynamical framework. The most convenient strategy for calculating the viscosity coefficients is to calculate the entropy generation in the CDM-radiation mixture within kinetic theory and equate it to the corresponding expression derived in the hydrodynamical framework. This allows to simply read off the viscosity coefficients by comparing the relevant tensors in both approaches that describe, respectively, the same type of entropy generation.

The effect of the viscous coupling is to damp perturbations in the CDM energy momentum tensor; in fact, they generate an exponential cut-off in the perturbation spectrum at kinetic decoupling. This spectrum then serves as initial condition for the subsequent evolution at temperatures $T < T_{\text{kd}}$ – the so-called free streaming regime, where a further damping in the perturbation spectrum takes place. The free streaming of CDM particles leaves a finger-print similar to collisional damping in the spectrum, i.e. a characteristic exponential damping term. Free streaming is therefore often also referred to as collision-less damping. For WIMPs, the characteristic damping mass is set by

$$\frac{M_{\text{fs}}}{10^{-6} M_\odot} \approx \left[\frac{1 + \ln(T_{\text{kd}}/30 \text{ MeV})/19.2}{(M_\chi/100 \text{ GeV})^{1/2} (T_{\text{kd}}/30 \text{ MeV})^{1/2}} \right]^3. \quad (27)$$

The characteristic damping scale for free streaming is time dependent, however, it becomes constant to a very good approximation sufficiently late after matter radiation equality.

The authors of [7, 8] calculated the envelope of the CDM perturbation spectrum. In [11] it was later shown numerically that acoustic oscillations have to be taken into account at kinetic decoupling as they give rise to even stronger damping; the order of magnitude of the cut-off scale, however, is correctly given in [7, 8]. Most recently, the author of [12] succeeded in including the acoustic oscillations before and during kinetic decoupling, the viscous coupling during kinetic decoupling and free streaming in a consistent kinetic treatment. All approaches agree in that the dominant suppression of small-scale power is given by collision-less damping during free streaming, with kinetic decoupling providing consistent initial conditions.

During radiation domination and on sub-Hubble scales, the transfer function for

matter density perturbations grows logarithmically, while after matter radiation equality matter density perturbations increase linearly (for a vanishing baryon fraction) with the scale factor on sub-Hubble scales. In both eras, during radiation and during matter domination, the spectrum of perturbations depends logarithmically on the comoving wavenumber that characterizes the corresponding mode. Hence, modes characterized by wavenumbers close to the cut-off in the power spectrum typically enter the nonlinear regime first. This is the basic picture of hierarchical structure formation: small scales collapse first to nonlinear structures, merging only later to larger structures. For WIMPs, this typically happens at a redshift between 40 and 80, depending on the interaction theory underlying the WIMP candidate and the details of the primordial power spectrum. All matter density perturbations that grow nonlinear are characterized by masses above M_{fs} .

Whether these first purely gravitationally bound WIMP structures can survive until today is under debate. Their mean density contrast is comparable to that of the galactic disk and an order of magnitude above that of the halo in the solar neighborhood. One has to keep in mind, however, that the very first perturbations entering the nonlinear regime are, rather, rare fluctuations with mass scales $M_{\text{rf}} > M_{\text{fs}}$. Those rare fluctuations can easily reach density contrasts which are an order of magnitude higher. High resolution simulations [26] show that structures with physical characteristics set by the cut-off in the linear power spectrum may survive merger processes of the first halos. The next critical phase starts with the formation of stars. Encounters with stars could disrupt the first CDM halos. So far, various studies [27, 28] show that the first halos would lose mass during an encounter, but their core should remain intact.

5. Impacts for the indirect detection of dark matter

Indirect detection methods for DM rely on the fact that most particle DM candidates can pair-annihilate into standard model particles. The annihilation products could then potentially be spotted in cosmic rays of various kinds, with an expected flux contribution that is proportional to the DM density squared. From hierarchical structure formation, as mentioned before, typical Milky-Way sized Halos are expected to contain a plethora of DM substructures rather than a homogeneous DM distribution. Focusing in the following on gamma rays, the consequences for indirect DM detection are two-fold and, as we will see, depend sensitively on an accurate determination of the cutoff scale in the power-spectrum as we have performed it in the preceding part of this work.

At this point, let us briefly introduce some necessary notation and denote by $\rho_{\text{sub}}(M_{\text{sub}}, \mathbf{r}_{\text{sub}})$ the density profile inside a typical dark matter subhalo (or “clump”) of mass M_{sub} (as opposed to the smooth, i.e., average DM density $\rho(\mathbf{r})$ in the Milky Way), and by $n_{\text{sub}}(M_{\text{sub}}, \mathbf{r})$ the number density of such clumps at a position $\mathbf{r}$ in the halo. For subhalos that cannot be resolved separately, the average annihilation flux

originating from $\mathbf{r}$ is then proportional to $\rho_{\text{eff}}^2(\mathbf{r})$, rather than $\rho^2(\mathbf{r})$, where

$$\rho_{\text{eff}}^2(\mathbf{r}) \equiv \int dM_{\text{sub}} \int d^3\mathbf{r}_{\text{sub}} n_{\text{sub}}(M_{\text{sub}}, \mathbf{r} - \mathbf{r}_{\text{sub}}) (\rho_{\text{sub}}(M_{\text{sub}}, \mathbf{r}_{\text{sub}}))^2 \quad (28)$$

$$\simeq \rho_0 \int dM_{\text{sub}} M_{\text{sub}} \delta(M_{\text{sub}}) n_{\text{sub}}(M_{\text{sub}}, \mathbf{r}). \quad (29)$$

The dimensionless quantity

$$\delta \equiv \frac{1}{\rho_0} \frac{\int d^3\mathbf{r}_{\text{sub}} (\rho_{\text{sub}}(M_{\text{sub}}, \mathbf{r}_{\text{sub}}))^2}{\int d^3\mathbf{r}_{\text{sub}} \rho_{\text{sub}}(M_{\text{sub}}, \mathbf{r}_{\text{sub}})} \quad (30)$$

that appears here is a measure of the effective density contrast between an average dark matter clump and the local halo density ρ_0 . The last step in (29) is valid if n_{sub} does not change very much on scales of the order of the size of a clump, i.e. for the whole region of integration one has $n_{\text{sub}}(M_{\text{sub}}, \mathbf{r} - \mathbf{r}_{\text{sub}}) \approx n_{\text{sub}}(M_{\text{sub}}, \mathbf{r})$.

Equipped with the above notations, we can now turn our discussion to the *total gamma-ray flux* from DM annihilations which is potentially strongly enhanced w.r.t. the case of a smooth halo distribution. To see this, note that the number density of subhalos scales with their mass as $n_{\text{sub}} \propto M_{\text{sub}}^{-2}$ – a behaviour that is expected from the theory of cosmological perturbations and has recently numerically been confirmed down to a mass scale of $4 \times 10^6 M_{\odot}$, corresponding to the resolution limit of state-of-the-art N -body simulations [14]. To a first approximation, the density contrast δ , on the other hand, scales only weakly with M_{sub} ; from (29), this means that the luminosity of individual subhalos is inversely proportional to M_{sub} , leading to an equal contribution to the total subhalo-induced gamma-ray flux from each decade in subhalo mass – which is precisely the behaviour that has numerically been found in [14]. As we have stressed before, it is therefore crucial to have an exact and reliable method to determine the cutoff in the spectrum below which there are no contributions to the annihilation flux. Extrapolating the results of the above-mentioned study down to the typical cutoff mass scales that we found earlier, $M_{\text{fs}} \sim 10^{-6} M_{\odot}$, we expect a factor of almost 4 for the increase of the expected annihilation signal as compared to that from a smooth halo distribution. Taking into account the effects of sub-subhalos may further enhance the annihilation flux significantly; on the other hand, reduced survival probabilities for very small clumps, as discussed in the previous Section, would lead to slightly less optimistic predictions. Each of these issues is far from being settled at the moment of writing and warrants more dedicated future studies on their own.

The other – perhaps even more important – effect of a large fraction of substructures in the DM halo distribution is the possibility of observing single DM clumps as gamma-ray point sources, maybe already in the near future with upcoming experiments like GLAST [29]. Comparing the projected all-sky GLAST sensitivity map for point-sources of DM annihilations [30] with the simulated gamma-ray sky of a Milky Way-like halo [14], unfortunately seems to suggest that, in order to actually resolve the brightest substructures, point-source sensitivities about two to three orders of magnitude better than the GLAST performance are needed. One has to keep in mind, however, that

the flux estimates obtained in [14] are extremely conservative in the sense that no substructures smaller than $\sim 10^4 M_\odot$ are resolved. This means, in particular, that the density profile *within* the clumps is essentially cut off at that scale; actual physical DM subhalos will exhibit much higher central densities and thus considerably enhanced fluxes (see also [14] for similar conclusions).

In order to quantify these last remarks, we adopt for the subhalo profiles the usual (α, β, γ) parametrization,

$$\rho_{\text{sub}}(\mathbf{r}) = \rho_{\text{sub}}(r_s) \left(\frac{r}{r_s} \right)^{-\gamma} \left[1 + \left(\frac{r}{r_s} \right)^\alpha \right]^{\frac{\gamma-\beta}{\alpha}}, \quad (31)$$

and assume it to be valid down to a scale $r_{\text{cut}} = \max\{r_{\text{kd}}, r_{\text{ann}}\}$ (here, r_{kd} corresponds to the earlier determined cut in the power spectrum due to damping effects after the kinetic decoupling of WIMPs from the heat bath, and r_{ann} is set by the maximal possible DM density due to DM self-annihilations [32]). The scale radius r_s determines the transition from large ($\rho_{\text{sub}} \propto r^{-\beta}$) to small ($\rho_{\text{sub}} \propto r^{-\gamma}$) radial distances; for the small subhalos we are interested in here, one expects rather large concentration parameters [26, 31], which means that the main contribution to the annihilation flux originates from the innermost regions of the subhalos and we can, for our purposes, well approximate the profile as $\rho_{\text{sub}} \propto r^{-\gamma}$. Since the annihilation flux from a single subhalo is proportional to δ , as defined in Eq.(30), we don't expect a strong dependence of this flux on the value of r_{cut} for the case of mildly cuspy profiles ($\gamma = 1$). For a more cuspy profile like $\gamma = 1.5$, on the other hand, we expect the annihilation flux to scale logarithmically with r_{cut} , thus leading to an enhancement of up to 30 w.r.t. to the flux considered in [14]. In fact, at least at higher redshifts, one expects even steeper profiles for small subhalos, with γ ranging from 1.5 to 2 [26], thus potentially further enhancing the resulting flux.

Of course, one has to keep in mind that neither the details of the innermost subhalo profiles nor the survival properties of these objects are sufficiently known at the present moment to make firm predictions. The numbers that we have presented above should therefore be considered as rather naive estimates, meant to illustrate the potential for GLAST – and certainly for successive, next-generation gamma-ray telescopes – to detect particularly bright subhalos as gamma-ray point sources. Note, in particular, that once GLAST has detected promising looking point sources, these could further be studied with air Cerenkov telescopes (ACTs) like CANGAROO [33], HESS [34], MAGIC [35] or VERITAS [36], taking advantage of the much better sensitivity of these telescopes as compared to satellite based experiments (up to three orders of magnitude at high energies – see, e.g. [6]).

6. Conclusions

The dominant damping effect for the evolution of linear CDM perturbations is given by collision-less damping due to the geodesic motion of CDM particles. Even though collisional damping is sub-dominant, it has to be taken into account during the kinetic

decoupling process, which provides consistent initial conditions for the free streaming regime. More importantly, collisional damping leads to a free streaming damping scale that depends on the temperature at kinetic decoupling. It is thus really this temperature scale, T_{kd} , that renders the position of the cut-off scale in the linear power spectrum sensitive to the WIMP micro-physics. Since the cut-off scale might have observable consequences on sub-pc scales, at least in principal, the kinetic decoupling history provides an exciting interface between the WIMP interaction theory and the substructures that are probed by direct and indirect dark matter searches.

We have presented, for the first time, a complete and analytic description of the kinetic decoupling process, without assuming a specific WIMP candidate; the master equation for the WIMP temperature evolution has been derived from first principles and accommodates any WIMP interaction theory. This could be achieved by separating slow and fast processes in the collision integral with a Born-Oppenheimer-type of ansatz, which proved to be an excellent approximation scheme for analyzing the WIMP radiation mixture – allowing, in particular, for a full error management; in this article, we have determined the decoupling scale to a precision of $p^2/M_\chi^2 \sim T_{\text{kd}}/M_\chi \sim 10^{-5}$.

The calculations presented here support the (order-of-magnitude) relaxation time approximation employed for the case of bino-like WIMPs before [9]; in this situation, we found a decoupling temperature that is about a factor of 2.7 smaller than the previous estimates, corresponding to a 4.2 times higher cutoff mass. As already mentioned, however, we were in particular able to significantly reduce the uncertainty in the determination of the cut-off scale. Let us stress again that this holds for *arbitrary* WIMP candidates and elastic scattering amplitudes. In particular, the method presented here can even be applied to regions of the parameter space where a simple scaling of the total cross-section with energy, e.g. $\sigma \sim \omega^0$ or $\sigma \sim \omega^2$ is no longer valid – as, e.g., in the case of *s*-channel resonances for LKP-like WIMPs. In this context, we would like to stress once more that it is, in fact, important to take into account the appropriate micro-physics; wrong assumptions about the particle nature of the DM candidate can lead to grossly wrong results in the determination of the cutoff scale.

Turning to explicit models, we have performed a complete scan over the mUED parameter space and found a very narrow range for the kinetic decoupling temperature: $27 \text{ MeV} \lesssim T_{\text{kd}} \lesssim 34 \text{ MeV}$. A parameter scan over the MSSM with the methods presented here lies beyond the scope of this work, however, see [18] for a current state of the art approach (using the order-of-magnitude relaxation time approximation). Another interesting direction for future studies would be to develop a framework for a kinetic description even for DM candidates like axions that never have been in thermal equilibrium.

Finally, we have commented on indirect detection prospects of a boosted signal from orthodox locations of enhanced dark matter densities, e.g. the galactic center or dwarf galaxies, and, much more promising, from individual dark matter clumps as gamma ray point sources. It remains, in fact, an exciting possibility that such objects may be seen already with the next generation of gamma-ray telescopes.

Acknowledgments

Research at Perimeter Institute for Theoretical Physics is supported in part by the Government of Canada through NSERC and by the Province of Ontario through MRI.

Appendix A. Boltzmann equation and temperature evolution in a Friedmann-Robertson-Walker spacetime

The Boltzmann equation is one of the fundamental equations of non-equilibrium statistical mechanics and describes the evolution of the phase space distribution function $f(\mathbf{x}, \mathbf{p}, t)$ for a given particle species χ . In an abstract form, it is usually written as

$$\hat{L}[f] = C[f]. \quad (\text{A.1})$$

Here, the Liouville operator $\hat{L}$ is the covariant generalization of the convective derivative familiar from hydrodynamics, or – in more technical terms – the variation with respect to an affine parameter λ along a geodesic:

$$\hat{L}[f] = \frac{df}{d\lambda} = \frac{dx^\mu}{d\lambda} \frac{\partial f}{\partial x^\mu} + \frac{dp^\mu}{d\lambda} \frac{\partial f}{\partial p^\mu} = p^\mu \frac{\partial f}{\partial x^\mu} - \Gamma_{\rho\sigma}^\mu p^\rho p^\sigma \frac{\partial f}{\partial p^\mu}. \quad (\text{A.2})$$

Since χ is supposed to be on shell, p^0 is actually not an independent degree of freedom – so in the above equation, we have $\partial f / \partial p^0 = 0$. Let us now consider a (flat) Friedmann-Robertson-Walker (FRW) spacetime,

$$ds^2 = dt^2 - a^2(t) d\mathbf{x}^2, \quad (\text{A.3})$$

and express everything in local (or *comoving*) momenta $\bar{\mathbf{p}} \equiv a\mathbf{p}$:

$$\hat{L}[f] = \frac{dx^\mu}{d\lambda} \frac{\partial f}{\partial x^\mu} + \frac{d\bar{p}^i}{d\lambda} \frac{\partial f}{\partial \bar{p}^i} \quad (\text{A.4})$$

$$= p^0 \left(\partial_t + a^{-1} \bar{\mathbf{p}} \cdot \nabla + \frac{da}{d\lambda} \bar{\mathbf{p}} \cdot \nabla_{\bar{\mathbf{p}}} - a \Gamma_{\rho\sigma}^i p^\rho p^\sigma \frac{\partial}{\partial \bar{p}^i} \right) f \quad (\text{A.5})$$

$$= p^0 (\partial_t - H \bar{\mathbf{p}} \cdot \nabla_{\bar{\mathbf{p}}}) f. \quad (\text{A.6})$$

In the last step, we have introduced $H \equiv \dot{a}/a$ and used the fact that the homogeneity of space implies $f = f(|\bar{\mathbf{p}}|, t)$. From now on, we will only use comoving momenta; for convenience, however, we will in the following drop the bars over the $\mathbf{p}$.

The collision term $C[f]$ on the right hand side of the Boltzmann equation takes account of all interactions that χ may experience. In Appendix B, we determine its exact analytical form for a non-relativistic particle that is in contact with a heat-bath of relativistic particles F through two-body scattering processes of the type $F\chi \rightarrow F\chi$. As explained in Section 2, this is precisely the situation that we encounter after the chemical decoupling of χ in the early universe. To leading order in T/M_χ and $\mathbf{p}^2/M_\chi^2$, where T is the temperature of the heat bath, the collision term then reads

$$C[f] = cM_\chi^2 \left(\frac{T}{M_\chi} \right)^{n+4} \{M_\chi T \Delta_{\mathbf{p}} + A \mathbf{p} \cdot \nabla_{\mathbf{p}} + B\} f(\mathbf{p}). \quad (\text{A.7})$$

In the above expression, n is determined by the low-energy behaviour of the scattering amplitude, see Eq. (B.21); c is an overall constant that is given in Eq. (B.22). Furthermore, we actually have $A = 1$ and $B = 3$, but in order to develop a physical understanding for the value of these constants, we will, for the time being, keep our discussion to the general form displayed in Eq. (A.7).

Let us start by dividing the Boltzmann equation by p^0 and then integrate it over all momenta. On the left-hand side, we find

$$\int \frac{d^3p}{(2\pi)^3} \{\partial_t - H\mathbf{p} \cdot \nabla_{\mathbf{p}}\} f(\mathbf{p}) = a^{-3} \partial_t (a^3 n_\chi), \quad (\text{A.8})$$

where n_χ is the number density of χ . Performing the same integral over (A.7) gives

$$\int \frac{d^3p}{(2\pi)^3} C[f]/p^0 = cM_\chi \left(\frac{T}{M_\chi}\right)^{n+4} (B - 3A) n_\chi. \quad (\text{A.9})$$

Of course, scattering processes should not change the comoving number density $a^3 n_\chi$ – which, from the above expressions, is only true for $B = 3A$.

Inspired by the situation for a thermal distribution, let us now *define* the “temperature” T_χ for an arbitrary distribution f of non-relativistic particles as

$$\int \frac{d^3p}{(2\pi)^3} \mathbf{p}^2 f(\mathbf{p}) \equiv 3M_\chi T_\chi n_\chi. \quad (\text{A.10})$$

With this definition, we have $T_\chi \approx T$ for a situation close to thermal equilibrium and $\epsilon \equiv (T - T_\chi)/T$ can therefore be considered as a parameter that characterizes the deviation of f from a thermal distribution. To determine the evolution of T_χ , we consider the next moment of the Boltzmann equation:

$$\int \frac{d^3p}{(2\pi)^3} \frac{\mathbf{p}^2}{p^0} \hat{L}[f] = \int \frac{d^3p}{(2\pi)^3} \frac{\mathbf{p}^2}{p^0} C[f]. \quad (\text{A.11})$$

Performing the integrals, and using $B = 3A$, we arrive at the following process equation:

$$T'_\chi - \left[2 + aA \left(\frac{T}{M_\chi}\right)^{n+2}\right] \frac{T_\chi}{T} = -a \left(\frac{T}{M_\chi}\right)^{n+2}, \quad (\text{A.12})$$

with

$$a \equiv \left(\frac{45}{\pi^3 g_{\text{eff}}}\right)^{1/2} \frac{M_{\text{Pl}}}{M_\chi} c. \quad (\text{A.13})$$

Here, $T'_\chi \equiv dT_\chi/dT$ and we have assumed that the universe is radiation dominated, with g_{eff} effective degrees of freedom in relativistic particles. For $n \neq -2$, Eq. (A.12) can be solved by the ansatz

$$T_\chi = \lambda(T) T_\chi^{(\text{hom})} = \frac{\lambda(T)}{M_\chi} T^2 \exp \left[\frac{aA}{n+2} \left(\frac{T}{M_\chi}\right)^{n+2} \right]. \quad (\text{A.14})$$

The general solution to the inhomogeneous differential equation (A.12) then reads:

$$T_\chi = \frac{T}{A} \left\{ 1 - \frac{z^{1/(n+2)}}{n+2} \exp[z] \Gamma[-(n+2)^{-1}, z] \right\}_{z=\frac{aA}{n+2} \left(\frac{T}{M_\chi}\right)^{n+2}} + \lambda_0 T_\chi^{(\text{hom})}. \quad (\text{A.15})$$




Figure A1. Phaseplot for the temperature evolution, expressed in the dimensionless quantities $x \equiv |a| (T/M_\chi)^{n+2}$ and $y \equiv T_\chi/T$. Here, we show the case for $n = 0$, but the same type of behaviour is obtained for any other (positive) value of n . As can be seen, any departure from thermal equilibrium ($T_\chi = T$) is restored almost immediately for $x \gtrsim 10$, forcing T_χ to follow the solid line which represents our general solution (A.22). For smaller temperatures, χ decouples from the heat bath; in this regime, its temperature is only influenced by the Hubble expansion and thus evolves as $T_\chi \propto T^2$.

In the limit of large and small T , respectively, the incomplete Gamma function behaves as

$$\Gamma[-(n+2)^{-1}, z] \xrightarrow{T \rightarrow \infty} z^{-1/(n+2)} \exp[-z]/z, \quad (\text{A.16})$$

$$\Gamma[-(n+2)^{-1}, z] \xrightarrow{T \rightarrow 0} \frac{n+2}{z^{1/(n+2)}} + \Gamma[-1/(n+2)]. \quad (\text{A.17})$$

Only for $A = 1$, as obtained in (B.22), one can thus recover the expected asymptotic behaviour (by choosing $\lambda_0 = 0$):

$$T_\chi \xrightarrow{T \rightarrow \infty} T, \quad (\text{A.18})$$

$$T_\chi \xrightarrow{T \rightarrow 0} \left(\frac{a}{n+2} \right)^{1/(n+2)} \Gamma \left[\frac{n+1}{n+2} \right] \frac{T^2}{M_\chi}. \quad (\text{A.19})$$

At high temperatures, the particles are still kept in thermodynamic equilibrium, i.e. $T_\chi = T$. The scaling $T_\chi \propto T^2$ at low temperatures, on the other hand, corresponds to the solution of the collision-less Boltzmann equation, i.e. Eq. (A.12) for $a = 0$.

To conclude our discussion of the process equation (A.12), let us rewrite it as

$$(n+2) \frac{dy}{dx} = \frac{y}{x} \pm (y-1), \quad (\text{A.20})$$

where we have introduced dimensionless variables

$$x \equiv |a| (T/M_\chi)^{n+2} \quad \text{and} \quad y \equiv T_\chi/T, \quad (\text{A.21})$$

and the upper (lower) sign applies if $a > 0$ ($a < 0$). In this form, we can immediately see that the asymptotic solution $y = 1$ for $x \ll 1$ (which corresponds to the expected $T_\chi = T$

for large T) is stable against small perturbations if and only if $a > 0$, i.e. $c > 0$. In fact, from the phaseplot shown in Fig. A1 it becomes apparent that in this case χ is kept in thermal equilibrium extremely efficiently for $x \gtrsim 10$ and that therefore the evolution of T_χ actually follows an attractor solution to Eq. (A.12). Once the temperature falls below $x \sim 10$, decoupling occurs on a rather short timescale and we observe the expected $T_\chi \propto T^2$ due to the Hubble expansion.

To summarize the results of this Appendix, we have developed a physical understanding for the functional form of the collision term (A.7) and found that T_χ evolves as a function of T according to

$$T_\chi = T \left\{ 1 - \frac{z^{1/(n+2)}}{n+2} \exp[z] \Gamma[-(n+2)^{-1}, z] \right\}_{z=\frac{a}{n+2} \left(\frac{T}{M_\chi}\right)^{n+2}}, \quad (\text{A.22})$$

essentially independent of the initial conditions. For convenience, we state here again the full expression for the constant a that appears above:

$$a \equiv \sum_f \left(\frac{10}{(2\pi)^9 g_{\text{eff}}} \right)^{1/2} g_{\text{SM}} c_n N_{n+3}^\pm \frac{M_{\text{Pl}}}{M_\chi}, \quad (\text{A.23})$$

where the sum runs over all relativistic scattering partners f (with g_{SM} internal degrees of freedom) and c_n is the corresponding scattering amplitude at zero momentum transfer, see Eq. (B.21); g_{eff} is the effective number of heat-bath degrees of freedom close to decoupling and $N_i^\pm$ is defined in Eqs. (B.23, B.24). Finally, we may now make use of the asymptotic behaviour, Eqs. (A.18, A.19), of our general solution to *define the kinetic decoupling temperature* by equating these two limits:

$$\frac{T_{\text{kd}}}{M_\chi} \equiv \left(\frac{T^2}{M_\chi T_\chi} \right)_{T \rightarrow 0} = \left(\left(\frac{a}{n+2} \right)^{1/(n+2)} \Gamma \left[\frac{n+1}{n+2} \right] \right)^{-1}. \quad (\text{A.24})$$

Appendix B. The collision term for two-body scattering processes

In this Appendix, we provide the details for the calculation of the collision term for scattering processes between a non-relativistic particle χ and relativistic (usually SM) heat-bath particles. For the former, we use $p^\mu = (E, \mathbf{p})$ to denote ingoing momenta, while for the latter we use $k^\mu = (\omega, \mathbf{k})$; the corresponding outgoing quantities are marked with a tilde. With these notations, the collision term takes the form

$$C = \int \frac{d^3 k}{(2\pi)^3 2\omega} \int \frac{d^3 \tilde{k}}{(2\pi)^3 2\tilde{\omega}} \int \frac{d^3 \tilde{p}}{(2\pi)^3 2\tilde{E}} (2\pi)^4 \delta^{(4)}(\tilde{p} + \tilde{k} - p - k) |\mathcal{M}|^2 J, \quad (\text{B.1})$$

where $\mathcal{M}$ is the scattering amplitude, summed over final and averaged over initial spin states, and

$$J \equiv g_{\text{SM}} [(1 \mp g^\pm)(\omega) g^\pm(\tilde{\omega}) f(\tilde{\mathbf{p}}) - (1 \mp g^\pm)(\tilde{\omega}) g^\pm(\omega) f(\mathbf{p})]. \quad (\text{B.2})$$

We will assume that the relativistic SM particles are thermally distributed, with g_{SM} being the number of spin states, so their phase-space number density is given by

$$g_{\text{SM}} g^\pm(\omega) = g_{\text{SM}} (e^{\omega/T} \pm 1)^{-1} \quad (\text{B.3})$$

for fermions (upper signs) and bosons (lower signs), respectively. The only assumption that we will make about the χ distribution function $f(\mathbf{p})$ is that we can neglect Pauli suppression factors – as we have done in (B.2). In the context that we are considering here, this is satisfied to a very good approximation.

Since, for kinematic reasons, the average momentum transferred during the scattering events is small, we can expand (B.1) as

$$C(E) = \sum_{j=0}^{\infty} C^j, \quad (\text{B.4})$$

$$C^j \equiv \int \frac{d^3k}{(2\pi)^3 2\omega} \int \frac{d^3\tilde{k}}{(2\pi)^3 2\tilde{\omega}} \int \frac{d^3\tilde{p}}{(2\pi)^3 2\tilde{E}} \times (2\pi)^4 \delta(\tilde{E} + \tilde{\omega} - E - \omega) |\mathcal{M}|^2 J \left[\frac{1}{j!} D_{\mathbf{q}}^j(\tilde{\mathbf{p}}) \delta^{(3)}(\tilde{\mathbf{p}} - \mathbf{p}) \right], \quad (\text{B.5})$$

where we have introduced

$$D_{\mathbf{q}}(\tilde{\mathbf{p}}) \equiv \mathbf{q} \cdot \nabla_{\tilde{\mathbf{p}}} \equiv (\tilde{\mathbf{k}} - \mathbf{k}) \cdot \nabla_{\tilde{\mathbf{p}}}, \quad (\text{B.6})$$

and the derivatives of the delta function that appear above are as usual defined in terms of integration by parts.

After these preliminaries, let us now start to calculate the expansion coefficients C^j . First, we note that $J|_{\tilde{\mathbf{p}}=\mathbf{p}, \tilde{\omega}=\omega} = 0$. From this, it immediately follows that

$$C^0 = 0. \quad (\text{B.7})$$

For the next coefficient, we find

$$C^1 = -2\pi \int \frac{d^3k}{(2\pi)^3 2\omega} \int \frac{d^3\tilde{k}}{(2\pi)^3 2\tilde{\omega}} \mathbf{q} \cdot \nabla_{\tilde{\mathbf{p}}} \left[\frac{|\mathcal{M}|^2 J}{2\tilde{E}} \delta(\tilde{E} + \tilde{\omega} - E - \omega) \right]_{\tilde{\mathbf{p}}=\mathbf{p}} \quad (\text{B.8})$$

$$= -\pi \int \frac{d^3k}{(2\pi)^3 2\omega} \int \frac{d^3\tilde{k}}{(2\pi)^3 2\tilde{\omega}} \times \delta(\tilde{\omega} - \omega) \frac{|\mathcal{M}|_{t=0}^2}{E} \left[\mathbf{q} \cdot \nabla_{\tilde{\mathbf{p}}} J - \frac{\mathbf{q} \cdot \mathbf{p}}{E} \partial_{\tilde{\omega}} J \right]_{\tilde{\mathbf{p}}=\mathbf{p}}, \quad (\text{B.9})$$

where we have used

$$\mathbf{q} \cdot \nabla_{\tilde{\mathbf{p}}} \delta(\tilde{E} + \tilde{\omega} - E - \omega) = \frac{\mathbf{q} \cdot \tilde{\mathbf{p}}}{\tilde{E}} \partial_{\tilde{\omega}} \delta(\tilde{E} + \tilde{\omega} - E - \omega) \quad (\text{B.10})$$

to perform the necessary integration by parts in $\tilde{\omega}$. Let us now write the scattering amplitude as a function of the Mandelstam variables s and t , where $s = M_\chi^2 + 2\omega(E - |\mathbf{p}| \cos \theta)$, and expand it in $|\mathbf{p}|/E$:

$$|\mathcal{M}|_{t=0}^2(s) = |\mathcal{M}|_{t=0, \theta=\pi/2}^2 - \omega \frac{|\mathbf{p}|}{E} \cos \theta \left(\partial_\omega |\mathcal{M}|_{t=0, \theta=\pi/2}^2 \right) + \mathcal{O}\left(\frac{\mathbf{p}^2}{E^2}\right). \quad (\text{B.11})$$

With $\mathbf{l} \parallel \mathbf{p} \parallel \nabla_{\tilde{\mathbf{p}}} J|_{\tilde{\mathbf{p}}=\mathbf{p}}$, the angular integrals that appear in the above expression for C^1 are then of the form

$$\int d\Omega \int d\tilde{\Omega} |\mathcal{M}|_{t=0}^2(\mathbf{q} \cdot \mathbf{l}) = \frac{16}{3} \pi^2 \omega^2 E^{-1} (\mathbf{p} \cdot \mathbf{l}) \left\{ \left(\partial_\omega |\mathcal{M}|_{t=0, \theta=\pi/2}^2 \right) + \mathcal{O}\left(\frac{\mathbf{p}^2}{E^2}\right) \right\}. \quad (\text{B.12})$$

To leading order in $\mathbf{p}^2/E^2$, and with the help of the useful relation

$$g^\pm(1 \mp g^\pm)(\omega) = \frac{e^{\omega/T}}{(e^{\omega/T} \pm 1)^2} = -T \partial_\omega g^\pm(\omega), \quad (\text{B.13})$$

we thus arrive at the following result for C^1 :

$$C^1 = \frac{g_{\text{SM}}}{6(2\pi)^3} M_\chi^{-3} \int d\omega \omega^4 (\partial_\omega g^\pm) \left(\partial_\omega |\mathcal{M}|_{\substack{t=0 \\ \theta=\pi/2}}^2 \right) [T M_\chi \mathbf{p} \cdot \nabla_{\mathbf{p}} + \mathbf{p}^2] f(\mathbf{p}). \quad (\text{B.14})$$

Let us now compute the next coefficient of the expansion (B.4):

$$C^2 = \pi \int \frac{d^3 k}{(2\pi)^3 2\omega} \int \frac{d^3 \tilde{k}}{(2\pi)^3 2\tilde{\omega}} (\mathbf{q} \cdot \nabla_{\tilde{\mathbf{p}}})^2 \left[\frac{|\mathcal{M}|^2 J}{2\tilde{E}} \delta(\tilde{E} + \tilde{\omega} - E - \omega) \right]_{\tilde{\mathbf{p}}=\mathbf{p}} \quad (\text{B.15})$$

$$\begin{aligned} &= \frac{\pi}{2} \int \frac{d^3 k}{(2\pi)^3 2\omega} \int \frac{d^3 \tilde{k}}{(2\pi)^3 2\tilde{\omega}} |\mathcal{M}|_{t=0}^2 \\ &\quad \times \left[\left(\frac{(\mathbf{q} \cdot \nabla_{\tilde{\mathbf{p}}})^2 J}{E} - 2 \frac{(\mathbf{q} \cdot \mathbf{p})(\mathbf{q} \cdot \nabla_{\tilde{\mathbf{p}}}) J}{E^3} \right) \delta(\tilde{\omega} - \omega) + \frac{(\mathbf{q} \cdot \mathbf{p})^2}{E^3} J \partial_{\tilde{\omega}}^2 \delta(\tilde{\omega} - \omega) \right. \\ &\quad \left. + \left(2 \frac{(\mathbf{q} \cdot \mathbf{p})(\mathbf{q} \cdot \nabla_{\tilde{\mathbf{p}}}) J}{E^2} + \frac{\mathbf{q}^2}{E^2} J - 3 \frac{(\mathbf{q} \cdot \mathbf{p})^2}{E^4} J \right) \partial_{\tilde{\omega}} \delta(\tilde{\omega} - \omega) \right]_{\tilde{\mathbf{p}}=\mathbf{p}} \quad (\text{B.16}) \end{aligned}$$

This time, the angular integrals are of the form

$$\int d\Omega \int d\tilde{\Omega} |\mathcal{M}|_{\substack{t=0 \\ \theta=\pi/2}}^2 q^i q^j = \frac{16}{3} \pi^2 (\omega^2 + \tilde{\omega}^2) |\mathcal{M}|_{\substack{t=0 \\ \theta=\pi/2}}^2 g^{ij} + \mathcal{O}\left(\frac{\mathbf{p}^2}{E^2}\right). \quad (\text{B.17})$$

Again, we will only keep the leading order terms and, after similar steps as for the computation of C^1 , we arrive at:

$$\begin{aligned} C^2 &= \frac{g_{\text{SM}}}{6(2\pi)^3} M_\chi^{-3} \int d\omega \omega^4 |\mathcal{M}|_{\substack{t=0 \\ \theta=\pi/2}}^2 \\ &\quad \times \left[-T M_\chi^2 \partial_\omega g^\pm \Delta_{\mathbf{p}} + M_\chi (4T \omega^{-1} \partial_\omega g^\pm - \partial_\omega g^\pm + T \partial_\omega^2 g^\pm) \mathbf{p} \cdot \nabla_{\mathbf{p}} \right. \\ &\quad \left. - 3M_\chi \partial_\omega g^\pm + \mathbf{p}^2 (4\omega^{-1} \partial_\omega g^\pm + \partial_\omega^2 g^\pm) \right] f(\mathbf{p}). \quad (\text{B.18}) \end{aligned}$$

The terms C^j for $j > 2$ only contribute higher-order corrections in $\mathbf{p}^2/E^2$ and ω/M_χ . To lowest non-vanishing order, and after the necessary partial integrations, we thus find the following expression for the collision integral:

$$C = C^1 + C^2 \quad (\text{B.19})$$

$$\begin{aligned} &= \frac{g_{\text{SM}}}{6(2\pi)^3} M_\chi^{-2} \int d\omega g^\pm(\omega) \partial_\omega \left(\omega^4 |\mathcal{M}|_{\substack{t=0 \\ s=M_\chi^2+2M_\chi\omega}}^2 \right) \\ &\quad \times \left[M_\chi T \Delta_{\mathbf{p}} + \mathbf{p} \cdot \nabla_{\mathbf{p}} + 3 \right] f(\mathbf{p}) \quad (\text{B.20}) \end{aligned}$$

Unless one encounters s -channel resonances (see Appendix C for a treatment of this case), one may now expand the amplitude at zero momentum transfer in ω/M_χ :

$$|\mathcal{M}|_{\substack{t=0 \\ s=M_\chi^2+2M_\chi\omega}}^2 \equiv c_n \left(\frac{\omega}{M_\chi} \right)^n + \mathcal{O}\left(\left(\frac{\omega}{M_\chi} \right)^{n+1} \right). \quad (\text{B.21})$$




Figure C1. Feynman diagrams for the elastic scattering of the LKP with SM fermions. In principle, there is also a contribution from a t -channel Higgs-exchange, but this is strongly suppressed due to the small Yukawa couplings involved.

Keeping only the lowest order term of this expansion, even the integral over the photon energies can be performed analytically and we arrive, finally, at the following expression:

$$C = \frac{g_{\text{SM}}}{6(2\pi)^3} M_\chi^2 c_n N_{n+3}^\pm \left(\frac{T}{M_\chi} \right)^{n+4} \left[M_\chi T \Delta_{\mathbf{p}} + \mathbf{p} \cdot \nabla_{\mathbf{p}} + 3 \right] f(\mathbf{p}), \quad (\text{B.22})$$

where (for $j > 0$)

$$N_j^+ \equiv \frac{j+1}{T^{j+1}} \int d\omega \omega^j g^+(\omega) = (1 - 2^{-j}) (j+1)! \zeta(j+1), \quad (\text{B.23})$$

$$N_j^- \equiv \frac{j+1}{T^{j+1}} \int d\omega \omega^j g^-(\omega) = (j+1)! \zeta(j+1). \quad (\text{B.24})$$

Note that in (B.22) – as in all corresponding formulas before – a summation over all SM scattering partners is understood.

The expression (B.22) is the central result of this Appendix; let us conclude by remarking that the formalism which we have presented here in some detail can straightforwardly be used to calculate even higher-order corrections to C . However, as we have learned in Appendix A, χ will be kept in thermal equilibrium until $\mathbf{p}^2/E^2, \langle \omega \rangle / M_\chi \sim 10^{-6} - 10^{-5}$, so the approximation to keep only leading order expressions in these quantities guarantees a result that is correct to a very high accuracy.

Appendix C. The LKP scattering amplitude

In this Appendix, we present the LKP scattering amplitude close to kinetic decoupling, which is dominated by the scattering with light SM fermions. The relevant Feynman diagrams are shown in Fig. C1. Note that in the UED model, at each given KK level, the number of fermions is doubled as compared to the SM, i.e. singlet and doublet states come each equipped with a whole KK tower of their own. The corresponding couplings of the LKP, well approximated by the $B^{(1)}$, are given by (see, e.g., [37]):

$$-g_Y \frac{Y_s}{2} B^{(1)}_\mu \bar{f}_s^{(1)} \gamma^\mu (1 + \gamma^5) f^{(0)} + c.c. \quad (\text{C.1})$$




Figure C2. The full LKP scattering amplitude (solid lines) vs. the approximative expression that scales with the photon energy as ω^2 (dashed lines). For simplicity, we have shown the case of electron scattering, and only taken into account the contribution from left-handed states. The two situations correspond to a mass splitting $\delta \equiv (M_{f_s^{(1)}} - M_{B(1)})/M_{B(1)}$ of 1% (left) and 5% (right), respectively.

for KK fermion singlet states, $f_s^{(1)}$, and

$$g_Y \frac{Y_d}{2} B_{(1)\mu} \bar{f}_d^{(1)} \gamma^\mu (1 - \gamma^5) f^{(0)} + c.c \quad (C.2)$$

for KK fermion doublet states, $f_d^{(1)}$. Given these couplings, and after summing over all initial states, it is straight-forward to calculate the amplitude at zero momentum transfer (in the limit of $m_f=0$):

$$|\mathcal{M}|_{t=0, s=M_\chi^2+2M_\chi\omega}^2 = \frac{16}{3} g_Y^4 Y_s^4 \omega^2 \frac{M_{B(1)}^2 \left(M_{f_s^{(1)}}^2 - M_{B(1)}^2 \right)^2 + 2\omega^2 \left(M_{f_s^{(1)}}^4 + 2M_{B(1)}^4 \right)}{\left[\left(M_{f_s^{(1)}}^2 - M_{B(1)}^2 \right)^2 - 4M_{B(1)}^2 \omega^2 \right]^2 + 16M_{B(1)}^2 M_{f_s^{(1)}}^2 \omega^2 \Gamma^2} + (s \leftrightarrow d), \quad (C.3)$$

where Γ is the decay width of the corresponding s -channel state. For fermion energies much smaller than the mass difference between the LKP and the KK leptons, $\omega/M_{B(1)} \ll \delta \equiv (M_{f^{(1)}} - M_{B(1)})/M_{B(1)}$, we may expand the scattering amplitude as we did in (B.21):

$$|\mathcal{M}|_{t=0, s=M_\chi^2+2M_\chi\omega}^2 \approx c_2 \left(\frac{\omega}{M_{B(1)}} \right)^2, \quad (C.4)$$

$$c_2 \equiv \frac{16}{3} g_Y^4 \left\{ Y_d^4 \left(\frac{M_{B(1)}^2}{M_{f_d^{(1)}}^2 - M_{B(1)}^2} \right)^2 + Y_s^4 \left(\frac{M_{B(1)}^2}{M_{f_s^{(1)}}^2 - M_{B(1)}^2} \right)^2 \right\}. \quad (C.5)$$

In Fig. C2, we compare the full expression for the scattering amplitude with the approximation of keeping only the lowest order term in $\omega/M_{B(1)}$; we can see that the

low-energy approximation is valid as long as $\omega \lesssim 0.1\delta M_{B(1)}$, and then starts to deviate considerably. Considering that we expect decoupling temperatures of $T_{\text{kd}} \sim 10 \text{ MeV}$ (see Section 3.3), this condition is easily satisfied for the *average* photon energy, $\langle\omega\rangle = 2.70T$, during the decoupling process. Still, one might worry about photons in the *tail* of the thermal distribution, which may very well hit the resonance and thereby enhance the effective scattering amplitude considerably.

To address this question, we have to go back to the full expression (B.20) for the collision integral that we derived in Appendix B. Going through similar steps as those leading to (A.12), we then find that the process equation for T_χ to be solved is

$$T'_\chi - [2 + a(T)] T_\chi/T = -a(T), \quad (\text{C.6})$$

where

$$a(T) = \sum_f \left(\frac{10}{(2\pi)^9 g_{\text{eff}}} \right)^{1/2} g_{\text{SM}} \frac{M_{\text{Pl}}}{M_{B(1)}^3} T^{-2} \int d\omega g^+(\omega) \partial_\omega \left(\omega^4 |\mathcal{M}|_{t=0}^2 \Big|_{s=M_\chi^2+2M_\chi\omega} \right) \quad (\text{C.7})$$

and the sum runs over all relativistic scattering partners. We may now compare $a(T)$, for $T \sim T_{\text{kd}}$, between the two situations where we either take the full expression (C.3) or the approximate expression (C.5) for the scattering amplitude. Using

$$\Gamma = \frac{g_Y^2}{8\pi} \delta^2 M_{B(1)}, \quad (\text{C.8})$$

which is valid in the limit of small mass splittings δ , we then find that for

$$\delta \gtrsim 3 \times 10^{-3} \quad (\text{C.9})$$

the mistake in $a(T)$ is less than 5% – which, from (A.24), roughly, corresponds to a 1% error in the estimate for T_{kd} . For smaller δ , on the other hand, the two ways to calculate $a(t)$ soon start to give grossly different results, rendering this type of error estimation invalid; in such a situation, one has to (numerically) study the asymptotic solution to the full process equation (C.6) in order to determine the decoupling scale. This shows that, indeed, the high-energy tail of the heat-bath fermions plays an important role in such a situation.

Recalling that in the mUED model we have $\delta \sim 10^{-2}$, we will here not further analyze the case of a very pronounced s -channel resonance. However, we note that mass splittings of the order of $\delta \sim 10^{-3}$ (and smaller) are actually feasible in, e.g., non-minimal UED versions. Here, we have provided all the necessary tools to correctly treat such situations.

References

- [1] D. N. Spergel *et al.*, arXiv:astro-ph/0603449.
- [2] D. J. Eisenstein *et al.* [SDSS Collaboration], *Astrophys. J.* **633**, 560 (2005) [arXiv:astro-ph/0501171].
- [3] S. Cole *et al.* [The 2dFGRS Collaboration], *Mon. Not. Roy. Astron. Soc.* **362** (2005) 505 [arXiv:astro-ph/0501174].

- [4] H. Hoekstra, H. K. C. Yee and M. D. Gladders, *Astrophys. J.* **577**, 595 (2002) [arXiv:astro-ph/0204295].
- [5] E. M. Leitch, J. M. Kovac, N. W. Halverson, J. E. Carlstrom, C. Pryke and M. W. E. Smith, *Astrophys. J.* **624**, 10 (2005) [arXiv:astro-ph/0409357].
- [6] G. Jungman, M. Kamionkowski and K. Griest, *Phys. Rept.* **267**, 195 (1996) [arXiv:hep-ph/9506380]. L. Bergström, *Rept. Prog. Phys.* **63**, 793 (2000) [arXiv:hep-ph/0002126]; G. Bertone, D. Hooper and J. Silk, *Phys. Rept.* **405**, 279 (2005) [arXiv:hep-ph/0404175].
- [7] A. M. Green, S. Hofmann and D. J. Schwarz, *Mon. Not. Roy. Astron. Soc.* **353**, L23 (2004) [arXiv:astro-ph/0309621].
- [8] A. M. Green, S. Hofmann and D. J. Schwarz, *JCAP* **0508**, 003 (2005) [arXiv:astro-ph/0503387].
- [9] S. Hofmann, D. J. Schwarz and H. Stoecker, *Phys. Rev. D* **64**, 083507 (2001) [arXiv:astro-ph/0104173].
- [10] V. Berezhinsky, V. Dokuchaev and Y. Eroshenko, *Phys. Rev. D* **68**, 103003 (2003) [arXiv:astro-ph/0301551].
- [11] A. Loeb and M. Zaldarriaga, *Phys. Rev. D* **71**, 103520 (2005) [arXiv:astro-ph/0504112].
- [12] E. Bertschinger, *Phys. Rev. D* **74**, 063509 (2006) [arXiv:astro-ph/0607319].
- [13] A. M. Green, S. Hofmann and D. J. Schwarz, *AIP Conf. Proc.* **805**, 431 (2006) [arXiv:astro-ph/0508553].
- [14] J. Diemand, M. Kuhlen and P. Madau, arXiv:astro-ph/0611370.
- [15] D. J. Fixsen, E. S. Cheng, J. M. Gales, J. C. Mather, R. A. Shafer and E. L. Wright, *Astrophys. J.* **473**, 576 (1996) [arXiv:astro-ph/9605054].
- [16] J. Bernstein, “KINETIC THEORY IN THE EXPANDING UNIVERSE,” Cambridge Univ. Press (1988).
- [17] A. S. Kompaneets, *Zh. Eksp. Teor. Fiz.* **31**, 876 (1957) [*Sov. Phys. JETP* **4**, 730 (1957)].
- [18] S. Profumo, K. Sigurdson and M. Kamionkowski, *Phys. Rev. Lett.* **97**, 031301 (2006) [arXiv:astro-ph/0603373].
- [19] T. Appelquist, H. C. Cheng and B. A. Dobrescu, *Phys. Rev. D* **64**, 035002 (2001) [arXiv:hep-ph/0012100].
- [20] H. C. Cheng, K. T. Matchev and M. Schmaltz, *Phys. Rev. D* **66**, 036005 (2002) [arXiv:hep-ph/0204342].
- [21] G. Servant and T. M. P. Tait, *Nucl. Phys. B* **650**, 391 (2003) [arXiv:hep-ph/0206071].
- [22] K. Kong and K. T. Matchev, *JHEP* **0601**, 038 (2006) [arXiv:hep-ph/0509119].
- [23] F. Burnell and G. D. Kribs, *Phys. Rev. D* **73**, 015001 (2006) [arXiv:hep-ph/0509118].
- [24] M. Kakizaki, S. Matsumoto and M. Senami, *Phys. Rev. D* **74**, 023504 (2006) [arXiv:hep-ph/0605280].
- [25] S. Weinberg, *Astrophys. J.* **168**, 175 (1971).
- [26] J. Diemand, B. Moore and J. Stadel, *Nature* **433**, 389 (2005) [arXiv:astro-ph/0501589].
- [27] A. M. Green and S. P. Goodwin, arXiv:astro-ph/0604142.
- [28] T. Goerdt, O. Y. Gnedin, B. Moore, J. Diemand and J. Stadel, arXiv:astro-ph/0608495.
- [29] <http://www-glast.stanford.edu/>
- [30] G. Bertone, T. Bringmann, R. Rando, G. Busetto and A. Morselli, arXiv:astro-ph/0612387.
- [31] P. Ullio, L. Bergstrom, J. Edsjo and C. G. Lacey, *Phys. Rev. D* **66**, 123502 (2002) [arXiv:astro-ph/0207125].
- [32] V. S. Berezhinsky, A. V. Gurevich and K. P. Zybin, *Phys. Lett. B* **294** (1992) 221.
- [33] <http://icrhp9.icrr.u-tokyo.ac.jp/index.html>
- [34] <http://www.mpi-hd.mpg.de/hfm/HESS/HESS.html>
- [35] <http://hegra1.mppmu.mpg.de/MAGICWeb/>
- [36] <http://veritas.sao.arizona.edu/index.html>
- [37] L. Bergström, T. Bringmann, M. Eriksson and M. Gustafsson, *JCAP* **0504**, 004 (2005) [arXiv:hep-ph/0412001].